

ProCHO References

ProCHO-4

1. Ambort D, Post S, Johansson M, et al. Function of the CysD domain of the gel-forming MUC2 mucin. *Biochem J.* 2011;436(Pt 1):61-70.
2. Austin A, Haas K, Naugler S, et al. Identification and Characterization of a Novel Regulatory Factor: IgA-Inducing Protein. *J Immunol.* 2003;171:1336-1342.
3. Austin A, Haas K, Naugler S, Bajer A, Garcia-Tapia D, Estes D. Identification and Characterization of a Novel Regulatory Factor: IgA-Inducing Protein. *The Journal of Immunology.* 2003;171(3):1336-1342.
4. Campo J, Lindqvist M, Cuello M, et al. Intranasal immunization with a proteoliposome-derived cochleate containing recombinant gD protein confers protective immunity against genital herpes in mice. *Vaccine.* 2010;28(5):1193-1200.
5. Domino S, Hurd E, Thomsson K, et al. Cervical mucins carry a(1,2) fucosylated glycans that partly protect from experimental vaginal candidiasis. *Glycoconjugate Journal.* 2009;26(9):1125-1134.
6. Egorova-Zachernyuk T, Bosman G, DeGrip W. Uniform stable-isotope labeling in mammalian cells: formulation of a cost-effective culture medium. *Applied Microbiology and Biotechnology.* 2011;89(2):397-406.
7. Freimark D, Jerome V, Freitag R. A GFP-based method facilitates clonal selection of transfected CHO cells. *Biotechnology Journal.* 2010;5(1):24-31.
8. Gao J, Cha S, Jonsson R, et al. Detection of anti-type 3 muscarinic acetylcholine receptor autoantibodies in the sera of Sjogren's syndrome patients by use of a transfected cell line assay. *Arthritis and Rheumatism.* 2004;50(8):2615-2621.
9. Joubert O, Nehme R, Fleury D, et al. Functional studies of membrane-bound and purified human Hedgehog receptor Patched expressed in yeast. *Biochimica et Biophysica Acta (BBA)- Biomembranes.* 2009;1788(9):1813-1821.
10. Klitgaard J, Coljee V, Andersen P, et al. Reduced Susceptibility of Recombinant Polyclonal Antibodies to Inhibitory Anti-Variable Domain Antibody Responses. *The Journal of Immunology.* 2006;177(6):3782-3790.
11. Link T, Backstrom M, Graham R, et al. Bioprocess development for the production of a recombinant MUC1 fusion protein expressed by CHO-K1 cells in a protein-free medium. *Journal of Biotechnology.* 2004;110(1):51-62.
12. Lutz S, Rao N, Wandrey C. Membranes in Biotechnology. *Chemical Engineering and Technology.* 2006;29(12):1404-1415.
13. Nielsen L, Baer A, Muller C, et al. Single-Batch Production of

- Recombinant Human Polyclonal Antibodies. *Molecular Biotechnology*. 2010;45(3):257-266.
14. Park J, Chio S, Park T. Enhancement of cell growth and viability of CHO cells in serum-free media by 30Kc6 gene expression. *Process Biochemistry*. 2007;42(1):8-15.
 15. Rasmussen L, Larsen Y, Hojrup P. Characterization of different cell culture media for expression of recombinant antibodies in mammalian cells: Presence of contaminating bovine antibodies. *Protein Expression and Purification*. 2005;41(2):373-377.
 16. Sewell R, Backstrom M, Dalziel et al. The ST6GalNAc-I Sialyltransferase Localizes throughout the Golgi and Is Responsible for the Synthesis of the Tumor-associated Sialyl-Tn O-Glycan in Human Breast Cancer. *The Journal of Biological Chemistry*. 2006;281:3586-3594.
 17. Sihlbom C, Hard I, Lidell M, et al. Localization of O-glycans in MUC1 glycoproteins using electron-capture dissociation fragmentation mass spectrometry. *Glycobiology*. 2009;19(4):375-381.
 18. Simon P, Brogle K, Wang B, Kyle D, Soltis D. Display of somatostatin-related peptides in the complementarity determining regions of an antibody light chain. *Archives of Biochemistry and Biophysics*. 2005;440(2):148-157.
 19. Singleton J, Guillen D, Scully M, et al. Characterization of Antibodies to CA 125 that Bind Preferentially to the Cell-Associated Form of the Antigen. *Tumor Biol*. 2006;27:122-132.
 20. Suck K, Walter J, Menzel F, et al. Fast and efficient protein purification using membrane adsorber systems. *Journal of Biotechnology*. 2006;121(3):361-367.
 21. Tsao M, Johnson J, Bachelder C, et al. Maintenance of High Viability During Direct Transition of CHO-K1 Cell Growth from 10% to 0% Serum; No Adaption Required. *Animal Cell Technology: Products From Cells, Cells As Products*. 2000;Chapter1;103-106.
 22. Walter J, Tappe A, Kasper C, et al. Human Growth Hormone (HGH) Purification From CHO-Cell Culture Supernatant Utilizing Macroporous Chromatographic Media. *Cell Technology for Cell Products*. 2007;Chapter VI:635-641.
 23. Wiendahl C, Brandner J, Klein B, et al. Metabolome Analysis in Mammalian Cells: Development and Application of a Sampling Technique. *ESACT Proceedings*. 2010;4(4):413-420.
 24. Wingens M, Gatgens J, Hoffrogge R, Noll T. Proteomic Characterization of a Glucose-Limited CHO Perfusion Process-Analysis of Metabolic Changes and Increase in Productivity. *ESACT Proceedings*. 2010;4(3):265-269.
 25. Ye J, Kober V, Tellers M, et al. High-level protein expression in scalable CHO transient transfection. *Biotechnology and Bioengineering*. 2009;103(3):542-551.
 26. Ziegler E, Gueler F, Rong S, et al. CCL19-IgG Prevents Allograft Rejection by Impairment of Immune Cell Trafficking. *JASN*. 2006;17(9):2521-2532.
 27. Ziegler E, Oberbarnscheidt M, Bulfone-Paus S, et al. CCR7 Signaling Inhibits T Cell Proliferation. *The Journal of Immunology*. 2007;179(10):6485-6493.

ProCHO-5

1. Abdulkerim E, Baganz S, Schambach A, Kasper C, Scheper T. Production and purification of TGFb-1 in CHO-Cells. *BMC Proc*. 2011;5(Suppl 8):P134.
2. Chenuet S, Martinet D, Besuchet-Schmutz N, et al. Calcium phosphate transfection generates mammalian recombinant cell lines with higher specific productivity than polyfection. *Biotechnology and Bioengineering*. 2008;101(5):937-945.
3. Croset A, Delafosse L, Gaudry J, et al. Differences in the glycosylation of

- recombinant proteins expressed in HEK and CHO cells. *Journal of Biotechnology*. 2012;161(3):336-348.
4. Derouazi M, Girard P, Tilborgh F, et al. Serum-free large-scale transient transfection of CHO cells. *Biotechnology and Bioengineering*. 2004;87(4):537-545.
 5. Duvar S, Berlin J, Ziehr H, Conradt H. Modulation of the Glycosylation Repertoire of a Recombinant Human EPO Expressing Model Cell Line Under Different Culture Conditions. *ESACT Proceedings*. 2005;2(5):723-725.
 6. Engelhardt E, Houis S, Gries T, Hilborn J, Adam M, Wurm F. Suspension-adapted Chinese hamster ovary-derived cells expressing green fluorescent protein as a screening tool for biomaterials. *Biotechnology Letters*. 2009;31(8):1143-1149.
 7. Freimark D, Jerome V, Freitag R, Effect of process parameters and product-host-interaction on hVEGFA-production by recombinant Chinese hamster ovary cells. *Biotechnology Progress*. 2012;28(3):762-772.
 8. Hacker D, Derow E, Wurm F. The CELO adenovirus Gam1 protein enhances transient and stable recombinant protein expression in Chinese hamster ovary cells. *The Journal of Biotechnology*. 2005;117(1):21-29.
 9. Marlind J, Kaspar M, Trachsel E, et al. Antibody-Mediated Delivery of Interleukin-2 to the Stroma of Breast Cancer Strongly Enhances the Potency of Chemotherapy. *Clinical Cancer Research*. 2008;14(20):6515-6524.
 10. Matasci M, Baldi L, Hacker D, Wurm F. The *PiggyBac* transposon enhances the frequency of CHO stable cell line generation and yields recombinant lines with superior productivity and stability. *Biotechnology and Bioengineering*. 2011;108(9):2141-2150.
 11. Muller N, Derouazi M, Tilborgh F, et al. Scalable transient gene expression in Chinese hamster ovary cells in instrumented and non-instrumented cultivation systems. *Biotechnology Letters*. 2007;29(5):703-711.
 12. Oberbek A, Matasci M, Hacker D, Wurm F. Generation of stable, high-producing CHO cell lines by lentiviral vector-mediated gene transfer in serum-free suspension culture. *Biotechnology and Bioengineering*. 2011;108(3):600-610.
 13. Rajendra Y, Backliwal G, Hildinger M, Wulhfard S, de Jesus M, Wurm F. A Transient Gene Expression Process with Recombinant Antibody Titer of 0.5g/L in CHO Cells. *ESACT Proceedings*. 2012;5(1):91-94.
 14. Rajendra Y, Kiseljak D, Manoli S, Baldi L, Hacker D, Wurm F. Role of non-specific DNA in reducing coding DNA requirement for transient gene expression with CHO and HEK-293E cells. *Biotechnology and Bioengineering*. 2012;109(9):2271-2278.
 15. Rajendra Y, Kiseljak D, Baldi L, Hacker D, Wurm F. A simple high-yielding process for transient gene expression in CHO cells. *Journal of Biotechnology*. 2011;153(1-2):22-26.
 16. Stettler M, Zhang X, Anderlei T, et al. Development of Pilot-Scale Orbital Shake Bioreactors: Ideal for Cost-Effective and Efficient Transient Gene Expression. *ESACT Proceedings*. 2010;4(3):183-185.
 17. Stettler M, Zhang X, Hacker D, de Jesus M, Wurm F. Novel Orbital Shake Bioreactors for Transient Production of CHO Derived IgGs. *Biotechnology Progress*. 2007;23(6):1340-1346.
 18. Wulhfard S, Tissot S, Bouchet S, et al. Mild Hypothermia Improves Transient Gene Expression Yields Several Fold in Chinese Hamster Ovary Cells. *Biotechnology Progress*. 2008;24(2):458-465.
 19. Wulhfard S, Baldi L, Hacker D, Wurm F. Valproic acid enhances recombinant mRNA and protein levels in transiently transfected Chinese hamster ovary cells. *Journal of Biotechnology*. 2010;148(2-3):128-132.
 20. Zhang X, Garcia I, Baldi L, Hacker D, Wurm F. Hyperosmolarity enhances transient recombinant protein yield in Chinese hamster ovary cells. *Biotechnology Letters*. 2010;32(11):1587-1592.
 21. Zhang X, Stettler M, Reif O, et al. Shaken helical track bioreactors: Providing oxygen

to high-density cultures of mammalian cells at volumes up to 1000L by surface aeration with air. *New Biotechnology*. 2008;25(1):68-75.