

MasterPure™ Yeast RNA Purification Kit

Cat. Nos. MPY03010 and MPY03100



Connect with Epicentre on our blog (epicentral.blogspot.com),
Facebook (facebook.com/EpicentreBio), and Twitter ([@EpicentreBio](https://twitter.com/EpicentreBio)).

1. Introduction

The MasterPure™ Yeast RNA Purification Kit provides all of the reagents needed to purify RNA from cell types including: *Candida*, *Saccharomyces*, *Schizosaccharomyces* and filamentous fungi. The kit utilizes a rapid desalting process¹ to remove contaminating macromolecules, avoiding toxic organic solvents, bead-beating, and spheroplasting.

2. Product Specifications

Storage: Store the Proteinase K and RNase-Free DNase I from the MasterPure Yeast RNA Purification Kit at –20°C in a freezer without a defrost cycle. The rest of the kit may be stored at room temperature for ease of use.

Storage Buffer: RNase-Free DNase I is supplied in a 50% glycerol solution containing 10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, and 10 mM CaCl₂. Proteinase K is supplied in a 50% glycerol solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 10 mM CaCl₂, 0.1% Triton® X-100, and 1 mM dithiothreitol (DTT).

Quality Control: The MasterPure Yeast RNA Purification Kit is function-tested by purifying RNA from *Saccharomyces cerevisiae*. RNA quality and yield are assayed by agarose gel electrophoresis, spectrophotometry and use as a template for RT-PCR.

3. Kit Contents

Desc.	Concentration	Quantity
The MasterPure Yeast RNA Purification Kit is available in 10- and 100-purification sizes. The 100-purification kit contains:		
Extraction Reagent for RNA		60 ml
MPC Protein Precipitation Reagent		50 ml
Proteinase K	@ 50 µg/µl	200 µl
RNase-Free DNase I	@ 1 U/µl	500 µl
2X T & C Lysis Solution		20 ml
RiboGuard™ RNase Inhibitor	@ 40 U/µl	100 µl
10X DNase Buffer	2 ml	
(330 mM Tris-HCl [pH 7.8], 660 mM potassium acetate, 100 mM magnesium acetate, and 5 mM DTT)		
TE Buffer		7 ml
(10 mM Tris-HCl [pH 7.5], 1 mM EDTA)		

4. Related Products

The following products are also available:

- MasterPure™ Yeast DNA Purification Kit

5. RNA Purification Protocol

A. RNA Purification

1. Dilute 1 µl of 50 µg/µl Proteinase K into 300 µl of Extraction Reagent for RNA for each sample. A premix may be prepared for multiple samples.
2. Pellet cells by centrifugation and discard the supernatant. The optimal number of cells varies with the species, but 1-1.5 ml of a mid-log culture gives good results for many types of yeast.
3. Vortex mix 10 seconds to loosen the cell pellet.
4. Add 300 µl of Extraction Reagent for RNA containing the Proteinase K and mix thoroughly by vortexing.
5. Incubate at 70°C for 10-15 minutes; vortex mix every 5 minutes. Shorter incubations of 5-10 minutes can yield nearly as much RNA.
6. Place the samples on ice for 3-5 minutes and add 175 µl of MPC Protein Precipitation Reagent to 300 µl of lysed sample (solution may become cloudy). Vortex vigorously for 10 seconds.
7. Pellet the debris by centrifugation for 10 minutes at 4°C at $\geq 10,000 \times g$ in a microcentrifuge.
8. Transfer the supernatant fluid to a clean microcentrifuge tube and discard the pellet.
9. Add 500 µl of isopropanol to the recovered supernatant fluid. Invert the tube 30-40 times.
10. Pellet the RNA by centrifugation at 4°C for 10 minutes at $\geq 10,000 \times g$ in a microcentrifuge.
11. Carefully pour off or aspirate the isopropanol without dislodging the RNA pellet. If removal of contaminating DNA is required, proceed with DNase I treatment in part B, otherwise, continue with this protocol. The MasterPure RNA Purification Kit extracts yeast DNA much less efficiently than RNA, and DNase I treatment may not be needed.
12. Rinse twice with 70% ethanol, being careful to not dislodge the pellet. Centrifuge briefly if the pellet is dislodged. Remove all of the residual ethanol with a pipet.
13. Resuspend the RNA in 35 µl of TE Buffer.
Note: TE Buffer is not supplied in the 10-purification size kit.
14. Quantitate RNA by its absorbance at 260 nm, and obtain an A_{260}/A_{280} ratio. Yields have been in the range of 25-50 µg of RNA per ml of $A_{600}=1.0$ cultures of *S. pombe*, and A_{260}/A_{280} ratios have been greater than 2.0.

B. Removal of Contaminating DNA from RNA Preparations

1. Remove all of the residual isopropanol with a pipet.
2. Prepare 200 µl of DNase I solution for each sample. Add 20 µl of 10X DNase Buffer to 175 µl of deionized water, then add 5 µl of RNase-Free DNase I.
3. Completely resuspend the nucleic acid pellet in 200 µl of DNase I solution.
4. Incubate at 37°C for 10 minutes.
Note: *Additional incubation (up to 30 min) may be necessary to remove all contaminating DNA.*
5. Add 200 µl of 2X T and C Lysis Solution; vortex mix for 5 seconds.
6. Add 200 µl of MPC Protein Precipitation Reagent (solution may become cloudy). Vortex mix 10 seconds; place on ice 3-5 minutes.
7. Pellet the debris by centrifugation at 4°C for 10 minutes at $\geq 10,000 \times g$ in a microcentrifuge.
8. Transfer the supernatant containing the RNA into a clean microcentrifuge tube and discard the pellet.
9. Add 500 µl of isopropanol to the supernatant. Invert the tube 30-40 times.
10. Pellet the purified RNA by centrifugation at 4°C for 10 minutes in a microcentrifuge at $\geq 10,000 \times g$.
11. Carefully pour off or aspirate the isopropanol without dislodging the RNA pellet.
12. Rinse twice with 70% ethanol, being careful to not dislodge the pellet. Centrifuge briefly if the pellet is dislodged. Remove all of the residual ethanol with a pipet.
13. Resuspend the RNA in 35 µl of TE Buffer.
14. Add 1 µl of RiboGuard™ RNase Inhibitor.

6. References

1. Miller, S.A. *et al.*, (1988) *Nucl. Acids Res.* **16**, 1215.

MasterPure and RiboGuard are trademarks of Epicentre, Madison, Wisconsin.

Triton is a registered trademark of Rohm & Haas, Philadelphia, Pennsylvania.

Visit our technical blog: epicentral.blogspot.com